arXiv:quant-ph/0210166v2 14 Dec 2002

N–Level System Interacting with Single Radiation Mode and Multi Cat States of Schrödinger in the Strong Coupling Regime

Kazuyuki FUJII *

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

In this paper we consider some (generalized) models which deal with n–level atom interacting with a single radiation mode and study a general structure of Rabi floppings of these generalized ones in the strong coupling regime.

To solve these models we introduce multi cat states of Schrödinger in our terminology and extend the result in (quant–ph/0203135) which Rabi frequencies are given by matrix elements of generalized coherent operators (quant–ph/0202081) under the rotating–wave approximation. Our work is a full generalization of Frasca (quant–ph/0111134) and Fujii (quant–ph/0203135).

In last we make a brief comment on an application to Quantum Computation on the space of qudits.

---

*E-mail address : fujii@yokohama-cu.ac.jp

1 Introduction

Coherent states or generalized coherent states play an important role in quantum physics, in particular, quantum optics, see [1] and [2]. They also play an important one in mathematical physics. See the textbook [3]. For example, they are very useful in performing stationary phase approximations to path integral, [4], [5], [6].

Coherent operators which produce coherent states are very useful because they are unitary and easy to handle. The basic reason is probably that they are subject to the elementary Baker-Campbell-Hausdorff (BCH) formula. Many basic properties of them are well–known, see [3] or [8].

Generalized coherent operators which produce generalized coherent states are also useful. But they are not so easy to handle in spite of having the disentangling one corresponding to the elementary BCH formula. In [7] and [16] the author determined all matrix elements of generalized coherent operators based on Lie algebras su(1,1) and su(2). They are interesting by themselves, but moreover have a very interesting application.

In [12] Frasca dealt with a model that describes a 2–level atom interacting with a single radiation mode (see [10] for a general review) in the strong coupling regime (not weak coupling one !) and showed that Rabi frequencies are obtained by matrix elements of coherent operator under the rotating–wave approximation. His aim was to explain the recent experimental finding on Josephson junctions [11].

This is an interesting result and moreover his method can be widely generalized. See also [13], [15] for another examples dealt with in the strong coupling regime.

His model has been generalized by Fujii [17] by making use of some operators based on Lie algebras su(1,1) and su(2), and a mathematical structure of Rabi floppings of these extended models in the strong coupling regime has been studied.

These models are based on 2–level atom interacting with a "single" radiation mode $\cdots$ 2–level system in the following. By the way, we are interested in n–level atom interacting with a "single" radiation mode $\cdots$ n–level system $\cdots$ which has been inspired by Quantum Computation, [20], [21], [22], [23], [24], [25]. That is, we would like to extend the results in 2–level system to ones in n–level system. This extension is not so easy.

In this paper we construct (propose) models that describe a n–level atom interacting with a "single" radiation mode. To solve these models we introduce multi cat states of Schrödinger (in our terminology) and show that (generalized) Rabi frequencies are also given by matrix elements of generalized coherent operators under the rotating–wave approximation.

We believe that the results will give a new aspect to Quantum Optics or Mathematical Physics.

Lastly we discuss an application to Quantum Computation on the space of qudits and discuss a possibility of application to Holonomic Quantum Computation, [26], [27], [28], [29], [30]–[34], but our discussion is not complete. See also Discussion in [17] concerning this possibility.

2 Coherent and Generalized Coherent Operators

2.1 Coherent Operator

Let $a(a^\dagger)$ be the annihilation (creation) operator of the harmonic oscillator. If we set $N \equiv a^\dagger a$ (: number operator), then

$$[N, a^\dagger] = a^\dagger \ , \ [N, a] = -a \ , \ [a^\dagger, a] = -\mathbf{1} \ . \tag{1}$$

Let $\mathcal{H}$ be a Fock space generated by a and $a^\dagger$, and $\{|n\rangle |\ n \in \mathbf{N} \cup \{0\}\}$ be its basis. The actions of a and $a^\dagger$ on $\mathcal{H}$ are given by

$$a|n\rangle = \sqrt{n}|n-1\rangle \ , \ a^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle \ , N|n\rangle = n|n\rangle \tag{2}$$

where $|0\rangle$ is a normalized vacuum ($a|0\rangle = 0$ and $\langle 0|0\rangle = 1$). From (2) state $|n\rangle$ for $n \geq 1$ are given by

$$|n\rangle = \frac{(a^\dagger)^n}{\sqrt{n!}}|0\rangle \ . \tag{3}$$

These states satisfy the orthogonality and completeness conditions

$$\langle m|n\rangle = \delta_{mn}\ , \quad \sum_{n=0}^{\infty} |n\rangle\langle n| = \mathbf{1}\ . \tag{4}$$

Definition We call a state

$$|z\rangle = \mathrm{e}^{za^{\dagger}-\bar{z}a}|0\rangle \equiv U(z)|0\rangle \quad \text{for} \quad z\ \in\ \mathbf{C} \tag{5}$$

the coherent state.

2.2 Generalized Coherent Operator Based on $su(1,1)$

Let us state generalized coherent operators and states based on $su(1,1)$.

We consider a spin K (>0) representation of $su(1,1) \subset sl(2,\mathbf{C})$ and set its generators $\{K_+, K_-, K_3\}$ $((K_+)^{\dagger} = K_-)$,

$$[K_3, K_+] = K_+, \quad [K_3, K_-] = -K_-, \quad [K_+, K_-] = -2K_3. \tag{6}$$

We note that this (unitary) representation is necessarily infinite dimensional. The Fock space on which $\{K_+, K_-, K_3\}$ act is $\mathcal{H}_K \equiv \{|K,n\rangle | n \in \mathbf{N}\cup\{0\}\}$ and whose actions are

$$\begin{aligned} &K_+|K,n\rangle = \sqrt{(n+1)(2K+n)}|K,n+1\rangle, \quad K_-|K,n\rangle = \sqrt{n(2K+n-1)}|K,n-1\rangle, \\ &K_3|K,n\rangle = (K+n)|K,n\rangle, \end{aligned} \tag{7}$$

where $|K,0\rangle$ is a normalized vacuum ($K_-|K,0\rangle = 0$ and $\langle K,0|K,0\rangle = 1$). We have written $|K,0\rangle$ instead of $|0\rangle$ to emphasize the spin K representation, see [4]. From (7), states $|K,n\rangle$ are given by

$$|K,n\rangle = \frac{(K_+)^n}{\sqrt{n!(2K)_n}}|K,0\rangle, \tag{8}$$

where $(a)_n$ is the Pochammer's notation $(a)_n \equiv a(a+1)\cdots(a+n-1)$. These states satisfy the orthogonality and completeness conditions

$$\langle K,m|K,n\rangle = \delta_{mn}, \quad \sum_{n=0}^{\infty} |K,n\rangle\langle K,n| \ = \mathbf{1}_K. \tag{9}$$

Now let us consider a generalized version of coherent states :

Definition We call a state

$$|z\rangle = V(z)|K,0\rangle \equiv \mathrm{e}^{zK_+ - \bar{z}K_-}|K,0\rangle \quad \text{for} \quad z \in \mathbf{C}. \tag{10}$$

the generalized coherent state (or the coherent state of Perelomov's type based on $su(1,1)$ in our terminology).

Here let us construct an example of this representation. First we set

$$K_+ \equiv \frac{1}{2}\left(a^\dagger\right)^2 \ , \ K_- \equiv \frac{1}{2}a^2 \ , \ K_3 \equiv \frac{1}{2}\left(a^\dagger a + \frac{1}{2}\right) \ , \tag{11}$$

then it is easy to check that these satisfy the commutation relations (6). That is, the set $\{K_+, K_-, K_3\}$ gives a unitary representation of $su(1,1)$ with spin $K = 1/4$ and $3/4$, [3]. Now we call an operator

$$S(z) = \mathrm{e}^{\frac{1}{2}\{z(a^\dagger)^2 - \bar{z}a^2\}} \quad \text{for} \quad z \in \mathbf{C} \tag{12}$$

the squeezed operator, see [3] or [19].

2.3 Generalized Coherent Operator Based on $su(2)$

Let us state generalized coherent operators and states based on $su(2)$.

We consider a spin J (> 0) representation of $su(2) \subset sl(2,\mathbf{C})$ and set its generators $\{J_+, J_-, J_3\}$ ($(J_+)^\dagger = J_-$),

$$[J_3, J_+] = J_+, \quad [J_3, J_-] = -J_-, \quad [J_+, J_-] = 2J_3. \tag{13}$$

We note that this (unitary) representation is necessarily finite dimensional. The Fock space on which $\{J_+, J_-, J_3\}$ act is $\mathcal{H}_J \equiv \{|J,n\rangle | 0 \le n \le 2J\}$ and whose actions are

$$\begin{aligned} &J_+|J,n\rangle = \sqrt{(n+1)(2J-n)}|J,n+1\rangle, \quad J_-|J,n\rangle = \sqrt{n(2J-n+1)}|J,n-1\rangle, \\ &J_3|J,n\rangle = (-J+n)|J,n\rangle, \end{aligned} \tag{14}$$

where $|J,0\rangle$ is a normalized vacuum ($J_-|J,0\rangle = 0$ and $\langle J,0|J,0\rangle = 1$). We have written $|J,0\rangle$ instead of $|0\rangle$ to emphasize the spin J representation, see [4]. From (14), states $|J,n\rangle$ are given by

$$|J,n\rangle = \frac{(J_+)^n}{\sqrt{n!_{2J}P_n}}|J,0\rangle. \tag{15}$$

These states satisfy the orthogonality and completeness conditions

$$\langle J,m|J,n\rangle=\delta_{mn},\quad \sum_{n=0}^{2J}|J,n\rangle\langle J,n|\;=\mathbf{1}_J. \tag{16}$$

Now let us consider a generalized version of coherent states :

Definition We call a state

$$|z\rangle=W(z)|J,0\rangle\equiv \mathrm{e}^{zJ_+-\bar{z}J_-}|J,0\rangle \quad \text{for} \quad z\in\mathbf{C}. \tag{17}$$

the generalized coherent state (or the coherent state of Perelomov's type based on $su(2)$ in our terminology).

2.4 Schwinger's Boson Method

Here let us construct the spin K and J representations by making use of Schwinger's boson method.

Next we consider the system of two-harmonic oscillators. If we set

$$a_1=a\otimes 1,\ {a_1}^\dagger=a^\dagger\otimes 1;\ a_2=1\otimes a,\ {a_2}^\dagger=1\otimes a^\dagger, \tag{18}$$

then it is easy to see

$$[a_i,a_j]=[{a_i}^\dagger,{a_j}^\dagger]=0,\ [a_i,{a_j}^\dagger]=\delta_{ij},\quad i,j=1,2. \tag{19}$$

We also denote by $N_i={a_i}^\dagger a_i$ number operators.

Now we can construct representation of Lie algebras $su(2)$ and $su(1,1)$ making use of Schwinger's boson method, see [4], [5]. Namely if we set

$$su(2):\quad J_+={a_1}^\dagger a_2,\ J_-={a_2}^\dagger a_1,\ J_3=\frac{1}{2}\left({a_1}^\dagger a_1-{a_2}^\dagger a_2\right), \tag{20}$$

$$su(1,1):\quad K_+={a_1}^\dagger {a_2}^\dagger,\ K_-=a_2a_1,\ K_3=\frac{1}{2}\left({a_1}^\dagger a_1+{a_2}^\dagger a_2+1\right), \tag{21}$$

then these satisfy the commutation relations (6) and (13) respectively.

3 Matrix Elements of Coherent and Generalized Coherent Operators $\cdots$ [16]

3.1 Matrix Elements of Coherent Operator

We list matrix elements of coherent operators $U(z)$.

The Matrix Elements The matrix elements of $U(z)$ are :

$$\text{(i)}\quad n \le m \quad \langle n|U(z)|m\rangle = \mathrm{e}^{-\frac{1}{2}|z|^2}\sqrt{\frac{n!}{m!}}(-\bar{z})^{m-n}{L_n}^{(m-n)}(|z|^2), \tag{22}$$

$$\text{(ii)}\quad n \ge m \quad \langle n|U(z)|m\rangle = \mathrm{e}^{-\frac{1}{2}|z|^2}\sqrt{\frac{m!}{n!}}z^{n-m}{L_m}^{(n-m)}(|z|^2), \tag{23}$$

where ${L_n}^{(\alpha)}$ is the associated Laguerre's polynomial defined by

$${L_k}^{(\alpha)}(x) = \sum_{j=0}^{k}(-1)^j\binom{k+\alpha}{k-j}\frac{x^j}{j!}. \tag{24}$$

In particular $L_k \equiv {L_k}^{(0)}$ is the usual Laguerre's polynomial and these are related to diagonal elements of $U(z)$.

3.2 Matrix Elements of Coherent Operator Based on $su(1,1)$

We list matrix elements of $V(z)$ coherent operators based on $su(1,1)$. In this case it is always $2K > 1$ ($2K = 1$ under some regularization).

The Matrix Elements The matrix elements of $V(z)$ are :

$$\begin{aligned}\text{(i)}\quad n \le m \quad \langle K,n|V(z)|K,m\rangle &= \sqrt{\frac{n!m!}{(2K)_n(2K)_m}}(-\bar{\kappa})^{m-n}(1+|\kappa|^2)^{-K-\frac{n+m}{2}} \times \\ &\sum_{j=0}^{n}(-1)^{n-j}\frac{\Gamma(2K+m+n-j)}{\Gamma(2K)(m-j)!(n-j)!j!}(1+|\kappa|^2)^j(|\kappa|^2)^{n-j},\end{aligned} \tag{25}$$

$$\begin{aligned}\text{(ii)}\quad n \ge m \quad \langle K,n|V(z)|K,m\rangle &= \sqrt{\frac{n!m!}{(2K)_n(2K)_m}}\kappa^{n-m}(1+|\kappa|^2)^{-K-\frac{n+m}{2}} \times \\ &\sum_{j=0}^{m}(-1)^{m-j}\frac{\Gamma(2K+m+n-j)}{\Gamma(2K)(m-j)!(n-j)!j!}(1+|\kappa|^2)^j(|\kappa|^2)^{m-j},\end{aligned} \tag{26}$$

where

$$\kappa \equiv \frac{\sinh(|z|)}{|z|}z = \cosh(|z|)\zeta. \tag{27}$$

The author doesn't know whether or not the right hand sides of (25) and (26) could be written by making use of some special functions such as generalized Laguerre's functions in (24). Therefore we set temporarily

$$F_m{}^{(n-m)}(x:2K)=\sum_{j=0}^{m}(-1)^{m-j}\frac{\Gamma(2K+m+n-j)}{\Gamma(2K)(m-j)!(n-j)!j!}(1+x)^j x^{m-j} \tag{28}$$

and $F_m{}^{(0)}(x;2K)=F_m(x;2K)$.

3.3 Matrix Elements of Coherent Operator Based on $su(2)$

We list matrix elements of $W(z)$ coherent operators based on $su(2)$. In this case it is always $2J\in\mathbf{N}$.

Matrix Elements The matrix elements of $W(z)$ are :

$$\text{(i)}\quad n\le m\quad \langle J,n|W(z)|J,m\rangle=\sqrt{\frac{n!m!}{{}_{2J}P_{n\,2J}P_m}}(-\bar{\kappa})^{m-n}(1-|\kappa|^2)^{J-\frac{n+m}{2}}\ \times$$
$$\sum_{j=0}^{n}{}_{*}(-1)^{n-j}\frac{(2J)!}{(2J-m-n+j)!(m-j)!(n-j)!j!}(1-|\kappa|^2)^j(|\kappa|^2)^{n-j}, \tag{29}$$

$$\text{(ii)}\quad n\ge m\quad \langle J,n|W(z)|J,m\rangle=\sqrt{\frac{n!m!}{{}_{2J}P_{n\,2J}P_m}}\kappa^{n-m}(1-|\kappa|^2)^{J-\frac{n+m}{2}}\ \times$$
$$\sum_{j=0}^{m}{}_{*}(-1)^{m-j}\frac{(2J)!}{(2J-m-n+j)!(m-j)!(n-j)!j!}(1-|\kappa|^2)^j(|\kappa|^2)^{m-j}, \tag{30}$$

where

$$\kappa\equiv\frac{\sin(|z|)}{|z|}z=\cos(|z|)\eta. \tag{31}$$

Here $\sum_{*}$ means a summation over j satisfying $2J-m-n+j\ge 0$.

The author doesn't know whether or not the right hand sides of (29) and (30) could be written by making use of some special functions. We set temporarily

$$F_m{}^{(n-m)}(x:2J)=\sum_{j=0}^{m}{}_{*}(-1)^{m-j}\frac{(2J)!}{(2J-m-n+j)!(m-j)!(n-j)!j!}(1-x)^j x^{m-j} \tag{32}$$

and $F_m{}^{(0)}(x;2J)=F_m(x;2J)$.

4 N–Level System in the Strong Coupling Region

Let us make a short review of [17] which is a generalization of [12].

In [12] Frasca dealt with a model that describes a 2–level atom interacting with a single radiation mode and developed some method to calculate Rabi frequencies in the strong coupling regime. We have generalized the model and method, and showed in [17] that Rabi frequencies in our extended model were given by matrix elements of generalized coherent operators ([16]) under the rotating–wave approximation.

Let $\{\sigma_1, \sigma_2, \sigma_3\}$ be Pauli matrices and $\mathbf{1}_2$ a unit matrix :

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{1}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{33}$$

We note here that

$$\sigma_1 = W\sigma_3 W^{-1} \tag{34}$$

where W is given by

$$W = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = W^{-1} \tag{35}$$

and is known as the Walsh–Hadamard matrix.

The Hamiltonians adopted in [17] are

$$\begin{aligned} &\text{(N)} \qquad H_N = \omega\mathbf{1}_2 \otimes a^\dagger a + \frac{\Delta}{2}\sigma_3 \otimes \mathbf{1} + g\sigma_1 \otimes (a^\dagger + a), \\ &\text{(K)} \qquad H_K = \omega\mathbf{1}_2 \otimes K_3 + \frac{\Delta}{2}\sigma_3 \otimes \mathbf{1}_K + g\sigma_1 \otimes (K_+ + K_-), \\ &\text{(J)} \qquad H_J = \omega\mathbf{1}_2 \otimes J_3 + \frac{\Delta}{2}\sigma_3 \otimes \mathbf{1}_J + g\sigma_1 \otimes (J_+ + J_-), \end{aligned} \tag{36}$$

where ω is the frequency of the radiation mode, Δ the separation between the two levels of the atom, g the coupling between the radiation field and the atom.

To deal with these three cases at the same time we set

$$\{L_+, L_-, L_3\} = \begin{cases} \text{(N)} & \{a^\dagger, a, N\} \\ \text{(K)} & \{K_+, K_-, K_3\} \\ \text{(J)} & \{J_+, J_-, J_3\} \end{cases} \tag{37}$$

and

$$H = H_0 + V = \omega\mathbf{1}_2 \otimes L_3 + \frac{\Delta}{2}\sigma_3 \otimes \mathbf{1}_L + g\sigma_1 \otimes (L_+ + L_-) \tag{38}$$

where we have written H instead of H_L for simplicity.

(36) is based on an atom with 2–level. However an atom has in reality many (energy) levels, so it is natural to consider an atom with n–level. First of all let us make some mathematical preliminaries, see for example [9]; Appendix.

Let $\{\Sigma_1, \Sigma_3\}$ be matrices in $M(n, \mathbf{C})$

$$\Sigma_1 = \begin{pmatrix} 0 & & & & & & 1 \\ 1 & 0 & & & & & \\ & 1 & 0 & & & & \\ & & 1 & \cdot & & & \\ & & & \cdot & \cdot & & \\ & & & & \cdot & \cdot & \\ & & & & & 1 & 0 \end{pmatrix}, \qquad \Sigma_3 = \begin{pmatrix} 1 & & & & & & \\ & \sigma & & & & & \\ & & \sigma^2 & & & & \\ & & & \cdot & & & \\ & & & & \cdot & & \\ & & & & & \cdot & \\ & & & & & & \sigma^{n-1} \end{pmatrix} \tag{39}$$

where σ is a primitive element satisfying $\sigma^n = 1$ ($\sigma = \mathrm{e}^{\frac{2\pi\sqrt{-1}}{n}}$). We note that $\{\Sigma_1, \Sigma_3\}$ is a generalization of Pauli matrices $\{\sigma_1, \sigma_3\}$, but they are not hermitian. If we define a Vandermonde matrix W based on σ as

$$W = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdots & \sigma^{(n-1)^2} \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdots & \sigma^{(n-1)(n-2)} \\ \cdot & \cdot & \cdot & & \cdot \\ \cdot & \cdot & \cdot & & \cdot \\ 1 & \sigma^2 & \sigma^4 & \cdots & \sigma^{2(n-1)} \\ 1 & \sigma & \sigma^2 & \cdots & \sigma^{n-1} \end{pmatrix}, \tag{40}$$

$$W^\dagger = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \sigma & \sigma^2 & \cdots & \sigma^{n-1} \\ 1 & \sigma^2 & \sigma^4 & \cdots & \sigma^{2(n-1)} \\ \cdot & \cdot & \cdot & & \cdot \\ \cdot & \cdot & \cdot & & \cdot \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdots & \sigma^{(n-1)(n-2)} \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdots & \sigma^{(n-1)^2} \end{pmatrix}, \tag{41}$$

then it is not difficult to see

$$\Sigma_1 = W\Sigma_3 W^\dagger = W\Sigma_3 W^{-1}. \tag{42}$$

For example, for $n = 3$

$$\begin{aligned} W\Sigma_3 W^\dagger &= \frac{1}{3}\begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} \begin{pmatrix} 1 & & \\ & \sigma & \\ & & \sigma^2 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix} \\ &= \frac{1}{3}\begin{pmatrix} 1 & \sigma & \sigma^2 \\ 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix} = \frac{1}{3}\begin{pmatrix} 0 & 0 & 3 \\ 3 & 0 & 0 \\ 0 & 3 & 0 \end{pmatrix} = \Sigma_1, \end{aligned}$$

where we have used that $\sigma^3 = 1,\ \bar{\sigma} = \sigma^2$ and $1 + \sigma + \sigma^2 = 0$.

That is, Σ_1 can be diagonalized by making use of W.

A comment is in order. Since W corresponds to the Walsh–Hadamard matrix (35), so it may be possible to call W the generalized Walsh–Hadamard matrix.

From this we know all eigenvalues and eigenstates of Σ_1 to become

$$\text{Eigenvalues} = \{1, \sigma, \cdots, \sigma^j, \cdots, \sigma^{n-1}\} \tag{43}$$

$$\text{Eigenstates} = \{|1\rangle, |\sigma\rangle, \cdots, |\sigma^j\rangle, \cdots, |\sigma^{n-1}\rangle\} \tag{44}$$

where@

$$|\sigma^j\rangle = \frac{1}{\sqrt{n}}\left(\begin{array}{c} 1 \\ \sigma^{(n-1)j} \\ \sigma^{(n-2)j} \\ \cdot \\ \cdot \\ \sigma^{2j} \\ \sigma^{j} \end{array}\right) \quad \text{for} \quad 0 \le j \le n-1.$$

Therefore Σ_1 can be written as (a spectral decomposition)

$$\Sigma_1 = \sum_{j=0}^{n-1} \sigma^j |\sigma^j\rangle\langle\sigma^j| \quad \text{and} \quad \mathbf{1}_n = \sum_{j=0}^{n-1} |\sigma^j\rangle\langle\sigma^j|. \tag{45}$$

The usual model is based on an atom with 2–level and we want to consider the case of an atom with n–level. Therefore it is very natural for us to generalize the model by making use of $\{\Sigma_1, \Sigma_3\}$ in place of $\{\sigma_1, \sigma_3\}$, but then we meet some trouble. One reason is Σ_1 and Σ_3 are **not** hermitian. The Hamiltonians must be hermitian in general.

After some trial and error we reach the following :

$$H = H_0 + V = \omega \mathbf{1}_n \otimes L_3 + \frac{\bar{\Delta}}{2}\Sigma_3 \otimes \mathbf{1}_L + \frac{\Delta}{2}{\Sigma_3}^{\dagger} \otimes \mathbf{1}_L + g(\Sigma_1 \otimes L_+ + {\Sigma_1}^{\dagger} \otimes L_-) \tag{46}$$

where Δ is a complex constant (we cannot prevent a **complex** constant). We believe that our extension is very natural except this point.

We cannot solve these simple models completely (maybe non–integrable), but we have found these models have a rich structure.

For these models we usually have two perturbation approaches :

Weak Coupling Regime $(0 < g \ll |\Delta|)$

$$H_0 = \omega \mathbf{1}_n \otimes L_3 + \frac{\bar{\Delta}}{2}\Sigma_3 \otimes \mathbf{1}_L + \frac{\Delta}{2}{\Sigma_3}^{\dagger} \otimes \mathbf{1}_L, \quad V = g(\Sigma_1 \otimes L_+ + {\Sigma_1}^{\dagger} \otimes L_-). \tag{47}$$

Strong Coupling Regime $(0 < |\Delta| \ll g)$

$$H_0 = \omega \mathbf{1}_n \otimes L_3 + g(\Sigma_1 \otimes L_+ + {\Sigma_1}^{\dagger} \otimes L_-), \quad V = \frac{\bar{\Delta}}{2}\Sigma_3 \otimes \mathbf{1}_L + \frac{\Delta}{2}{\Sigma_3}^{\dagger} \otimes \mathbf{1}_L. \tag{48}$$

In this paper we are not interested in the weak coupling regime, so in the following we focus our attention on the strong coupling regime. We would like to generalize the result in [17].

Then we have from (45) and (48)

$$\begin{aligned}
H_0 &= \sum_{j=0}^{n-1} |\sigma^j\rangle\langle\sigma^j| \otimes \{\omega L_3 + g(\sigma^j L_+ + \bar{\sigma}^j L_-)\} \\
&= \sum_{j=0}^{n-1} |\sigma^j\rangle\langle\sigma^j| \otimes \left\{\mathrm{e}^{-\frac{1}{2}(z_j L_+ - \bar{z}_j L_-)}\,(\Omega L_3)\,\mathrm{e}^{\frac{1}{2}(z_j L_+ - \bar{z}_j L_-)}\right\} \\
&= \sum_{j=0}^{n-1} \left(|\sigma^j\rangle \otimes \mathrm{e}^{-\frac{1}{2}(z_j L_+ - \bar{z}_j L_-)}\right)(\Omega L_3)\left(\langle\sigma^j| \otimes \mathrm{e}^{\frac{1}{2}(z_j L_+ - \bar{z}_j L_-)}\right) \qquad (49)
\end{aligned}$$

where we have used the following

Key Formulas

$$(N)\quad \omega a^\dagger a + g(\sigma^j a^\dagger + \bar{\sigma}^j a) = \Omega\ \mathrm{e}^{-\frac{1}{2}(z_j a^\dagger - \bar{z}_j a)}\left(N - \frac{g^2}{\omega^2}\right)\mathrm{e}^{\frac{1}{2}(z_j a^\dagger - \bar{z}_j a)}$$

$$\text{where}\quad \Omega = \omega, \quad z_j = (2g/\omega)\sigma^j, \qquad (50)$$

$$(K)\quad \omega K_3 + g(\sigma^j K_+ + \bar{\sigma}^j K_-) = \Omega\ \mathrm{e}^{-\frac{1}{2}(z_j K_+ - \bar{z}_j K_-)} K_3\ \mathrm{e}^{\frac{1}{2}(z_j K_+ - \bar{z}_j K_-)}$$

$$\text{where}\quad \Omega = \omega\sqrt{1-(2g/\omega)^2}, \quad z_j = \tanh^{-1}(2g/\omega)\sigma^j, \qquad (51)$$

$$(J)\quad \omega J_3 + g(\sigma^j J_+ + \bar{\sigma}^j J_-) = \Omega\ \mathrm{e}^{-\frac{1}{2}(z_j J_+ - \bar{z}_j J_-)} J_3\ \mathrm{e}^{\frac{1}{2}(z_j J_+ - \bar{z}_j J_-)}$$

$$\text{where}\quad \Omega = \omega\sqrt{1+(2g/\omega)^2}, \quad z_j = \tan^{-1}(2g/\omega)\sigma^j. \qquad (52)$$

These formulas are a slight generalization of corresponding ones in [17]. We leave the proof to the readers.

That is, we could diagonalize the Hamiltonian H_0. This is n–fold degenerate and its eigenvalues and eigenvectors are given respectively

$$(\text{Eigenvalues}, \text{Eigenvectors}) = \begin{cases} (N) \quad \Omega(-\frac{g^2}{\omega^2} + m), \quad |\sigma^j\rangle \otimes \mathrm{e}^{-\frac{1}{2}(z_j a^\dagger - \bar{z}_j a)}|m\rangle \\ (K) \quad \Omega(K + m), \quad |\sigma^j\rangle \otimes \mathrm{e}^{-\frac{1}{2}(z_j K_+ - \bar{z}_j K_-)}|K, m\rangle \\ (J) \quad \Omega(-J + m), \quad |\sigma^j\rangle \otimes \mathrm{e}^{-\frac{1}{2}(z_j J_+ - \bar{z}_j J_-)}|J, m\rangle \end{cases} \qquad (53)$$

for $j = 0, 1, \cdots, n-1$ and $m \in \mathbf{N} \cup \{0\}$. For the latter convenience we set

$$\text{Eigenvalues} = \{E_m\}, \quad \text{Eigenvectors} = \{|\{\sigma^j, m\}\rangle\}. \qquad (54)$$

Then (49) can be written as

$$H_0 = \sum_j \sum_m E_m |\{\sigma^j, m\}\rangle\langle\{\sigma^j, m\}|. \tag{55}$$

Next we would like to solve the following Schrödinger equation :

$$i\frac{d}{dt}\Psi = H\Psi = \left(H_0 + \frac{\bar{\Delta}}{2}\Sigma_3 \otimes \mathbf{1}_L + \frac{\Delta}{2}{\Sigma_3}^\dagger \otimes \mathbf{1}_L\right)\Psi, \tag{56}$$

where we have set $\hbar = 1$ for simplicity. To solve this equation we appeal to the **method of constant variation**. First let us solve

$$i\frac{d}{dt}\Psi = H_0\Psi, \tag{57}$$

which general solution is given by

$$\Psi(t) = U_0(t)\Psi_0 = \mathrm{e}^{-itH_0}\Psi_0 \tag{58}$$

where Ψ_0 is a constant state. It is easy to see from (55)

$$U_0(t) = \mathrm{e}^{-itH_0} = \sum_j \sum_m \mathrm{e}^{-itE_m} |\{\sigma^j, m\}\rangle\langle\{\sigma^j, m\}|. \tag{59}$$

The method of constant variation goes as follows. Changing like $\Psi_0 \longrightarrow \Psi_0(t)$, we insert (58) into (56). After some algebra we obtain

$$i\frac{d}{dt}\Psi_0 = \left\{\frac{\Delta}{2}{U_0}^\dagger\left({\Sigma_3}^\dagger \otimes \mathbf{1}_L\right)U_0 \ + \ \frac{\bar{\Delta}}{2}{U_0}^\dagger\left(\Sigma_3 \otimes \mathbf{1}_L\right)U_0\right\}\Psi_0 \equiv (H_F + {H_F}^\dagger)\Psi_0. \tag{60}$$

Now let us calculate H_F. From (59) and (53)

$$\begin{aligned} H_F &= \frac{\Delta}{2}\sum_{j,k}\sum_{m,r} \mathrm{e}^{it(E_m - E_r)}\langle\{\sigma^j, m\}|({\Sigma_3}^\dagger \otimes \mathbf{1}_L)|\{\sigma^k, r\}\rangle \ |\{\sigma^j, m\}\rangle\langle\{\sigma^k, r\}| \\ &= \frac{\Delta}{2}\sum_j \sum_{m,r} \mathrm{e}^{it\Omega(m-r)}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_j - z_{j-1})L_+ - (\bar{z}_j - \bar{z}_{j-1})L_-\}}|r\rangle\rangle \ |\{\sigma^j, m\}\rangle\langle\{\sigma^{j-1}, r\}| \end{aligned} \tag{61}$$

where we have used the relation $\langle\sigma^j|{\Sigma_3}^\dagger = \langle\sigma^{j-1}|$. Remind that $|m\rangle\rangle$ is respectively

$$|m\rangle\rangle = \begin{cases} (N) & |m\rangle \\ (K) & |K, m\rangle \\ (J) & |J, m\rangle \end{cases} \tag{62}$$

In this stage we meet **matrix elements of the coherent and generalized coherent operators** $\mathrm{e}^{zL_+-\bar{z}L_-}$ in section 3. Here we divide H_F into two parts

$$H_F = {H_F}^{'} + {H_F}^{''}$$

where

$${H_F}^{'} = \frac{\Delta}{2}\sum_j \sum_m \langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_j-z_{j-1})L_+-(\bar{z}_j-\bar{z}_{j-1})L_-\}}|m\rangle\rangle\ |\{\sigma^j,m\}\rangle\langle\{\sigma^{j-1},m\}|, \tag{63}$$

$${H_F}^{''} = \frac{\Delta}{2}\sum_j \sum_{\substack{m,r\\ m\neq r}} \mathrm{e}^{it\Omega(m-r)}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_j-z_{j-1})L_+-(\bar{z}_j-\bar{z}_{j-1})L_-\}}|r\rangle\rangle\ |\{\sigma^j,m\}\rangle\langle\{\sigma^{j-1},r\}|. \tag{64}$$

Here let us deal with ${H_F}^{'}$. By the formulas in section 3 it is easy to see

$$\begin{aligned}&\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_0-z_{-1})L_+-(\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle = \langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_1-z_0)L_+-(\bar{z}_1-\bar{z}_0)L_-\}}|m\rangle\rangle = \cdots\\ =&\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_{n-1}-z_{n-2})L_+-(\bar{z}_{n-1}-\bar{z}_{n-2})L_-\}}|m\rangle\rangle\end{aligned} \tag{65}$$

because by **Key Formulas** ((50), (51), (52))

$$z_j - z_{j-1} = C\sigma^{j-1}(\sigma-1) = C\sigma^{j-1}\sigma^{1/2}(\sigma^{1/2}-\sigma^{-1/2}) = 2C\mathrm{sin}(\pi/n)i\sigma^{j-1/2}, \tag{66}$$

$$|z_j - z_{j-1}| = 2C\mathrm{sin}(\pi/n) \quad \text{for} \quad j = 0,1,\cdots,n-1$$

where C is a constant defined by

$$C = \begin{cases} (N) & 2g/\omega \\ (K) & \tanh^{-1}(2g/\omega) \\ (J) & \tan^{-1}(2g/\omega) \end{cases} \tag{67}$$

Therefore ${H_F}^{'}$ can be written as

$${H_F}^{'} = \frac{\Delta}{2}\sum_m \langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_0-z_{-1})L_+-(\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle \left\{\sum_j |\{\sigma^j,m\}\rangle\langle\{\sigma^{j-1},m\}|\right\}.$$

Let us diagonalize the last term. For that we set for simplicity

$$\Lambda_m = \Big(|\{1,m\}\rangle, |\{\sigma^1,m\}\rangle, \cdots, |\{\sigma^{n-1},m\}\rangle\Big).$$

Then it is easy to see from (42)

$$\sum_{j=0}^{n-1} |\{\sigma^j,m\}\rangle\langle\{\sigma^{j-1},m\}| = \Lambda_m\Sigma_1\Lambda_m^\dagger = \Lambda_m W\Sigma_3 W^\dagger\Lambda_m^\dagger \equiv \widetilde{\Lambda}_m\Sigma_3\widetilde{\Lambda}_m^\dagger.$$

Here if we introduce a new basis

$$\Lambda_m W = \widetilde{\Lambda}_m = \left(|\{1,\psi_m\}\rangle, |\{\sigma,\psi_m\}\rangle, \cdots, |\{\sigma^{n-1},\psi_m\}\rangle \right),$$

where

$$\begin{aligned} |\{\sigma^j,\psi_m\}\rangle &\equiv (\widetilde{\Lambda}_m)_j = \sum_{k=0}^{n-1} (\Lambda_m)_k W_{kj} \\ &= \frac{1}{\sqrt{n}} \left(|\{1,m\}\rangle + \sigma^{j(n-1)}|\{\sigma^1,m\}\rangle + \sigma^{j(n-2)}|\{\sigma^2,m\}\rangle + \cdots + \sigma^j|\{\sigma^{n-1},m\}\rangle \right) \end{aligned} \tag{68}$$

then we obtain

$$H_F{}' = \frac{\Delta}{2} \sum_m \sum_j \langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_0-z_{-1})L_+ - (\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle \sigma^j |\{\sigma^j,\psi_m\}\rangle\langle\{\sigma^j,\psi_m\}|. \tag{69}$$

This is just diagonal !

A comment is in order. We would like to call the states (68) **multi cat states of Schrödinger** following from [19] (because we are dealing with multi level systems).

Next let us deal with $H_F{}''$. By $\Lambda_m = \widetilde{\Lambda}_m W^\dagger$ and (40), (41) we have

$$\begin{aligned} &|\{\sigma^j,m\}\rangle\langle\{\sigma^{j-1},r\}| = (\Lambda_m)_j ({\Lambda_r}^\dagger)_{j-1} = (\widetilde{\Lambda}_m W^\dagger)_j (W \widetilde{\Lambda}_r^\dagger)_{j-1} \\ &= \sum_{k=0}^{n-1}\sum_{l=0}^{n-1} (\widetilde{\Lambda}_m)_k (W^\dagger)_{kj} W_{j-1,l} (\widetilde{\Lambda}_r^\dagger)_l = \sum_{k,l} (W^\dagger)_{kj} W_{j-1,l} |\{\sigma^k,\psi_m\}\rangle\langle\{\sigma^l,\psi_r\}| \\ &= \frac{1}{n}\sum_{k,l} \sigma^{jk+(n-j+1)l} |\{\sigma^k,\psi_m\}\rangle\langle\{\sigma^l,\psi_r\}| \quad \text{or} \quad = \frac{1}{n}\sum_{k,l} \sigma^{jk-(j-1)l} |\{\sigma^k,\psi_m\}\rangle\langle\{\sigma^l,\psi_r\}|, \end{aligned}$$

so that we obtain

$$\begin{aligned} H_F{}'' = \frac{\Delta}{2} \sum_{\substack{m,r \\ m\neq r}} \sum_{k,l} \mathrm{e}^{it\Omega(m-r)} &\left\{ \sum_{j=0}^{n-1} \langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_j-z_{j-1})L_+ - (\bar{z}_j-\bar{z}_{j-1})L_-\}}|r\rangle\rangle \right. \times \\ &\left. \frac{\sigma^{jk-(j-1)l}}{n} |\{\sigma^k,\psi_m\}\rangle\langle\{\sigma^l,\psi_r\}| \right\}. \end{aligned} \tag{70}$$

Let us sum up the above result :

$$H_F = H_F{}' + H_F{}'', \quad H_F{}' = (69) \quad \text{and} \quad H_F{}'' = (70).$$

Next we take the dagger of H_F which is now an easy task.

$$H_F{}^\dagger \equiv \tilde{H}_F' + \tilde{H}_F''$$

where

$$\tilde{H}_F^{'} = \frac{\bar{\Delta}}{2}\sum_m \sum_j \langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_0-z_{-1})L_+-(\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle \sigma^{-j}|\{\sigma^j,\psi_m\}\rangle\langle\{\sigma^j,\psi_m\}|, \tag{71}$$

and

$$\begin{aligned}
\tilde{H}_F^{''} &= \frac{\bar{\Delta}}{2}\sum_{\substack{m,r\\ m\neq r}}\sum_{k,l}\mathrm{e}^{-it\Omega(m-r)}\left\{\sum_{j=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_j-z_{j-1})L_+-(\bar{z}_j-\bar{z}_{j-1})L_-\}}|m\rangle\rangle \quad\times\right.\\
&\qquad\left.\frac{\bar{\sigma}^{jk-(j-1)l}}{n}|\{\sigma^l,\psi_r\}\rangle\langle\{\sigma^k,\psi_m\}|\right\}\\
&= \frac{\bar{\Delta}}{2}\sum_{\substack{m,r\\ m\neq r}}\sum_{k,l}\mathrm{e}^{it\Omega(r-m)}\left\{\sum_{j=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_j-z_{j-1})L_+-(\bar{z}_j-\bar{z}_{j-1})L_-\}}|m\rangle\rangle \quad\times\right.\\
&\qquad\left.\frac{\sigma^{(j-1)l-jk}}{n}|\{\sigma^l,\psi_r\}\rangle\langle\{\sigma^k,\psi_m\}|\right\}\\
&= \frac{\bar{\Delta}}{2}\sum_{\substack{m,r\\ m\neq r}}\sum_{k,l}\mathrm{e}^{it\Omega(m-r)}\left\{\sum_{j=0}^{n-1}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_j-z_{j-1})L_+-(\bar{z}_j-\bar{z}_{j-1})L_-\}}|r\rangle\rangle \quad\times\right.\\
&\qquad\left.\frac{\sigma^{(j-1)k-jl}}{n}|\{\sigma^k,\psi_m\}\rangle\langle\{\sigma^l,\psi_r\}|\right\}.
\end{aligned} \tag{72}$$

Therefore

$$H_F + {H_F}^{\dagger} = \left({H_F}^{'} + \tilde{H}_F^{'}\right) + \left({H_F}^{''} + \tilde{H}_F^{''}\right),$$

where

$${H_F}^{'} + \tilde{H}_F^{'} = \sum_m\sum_j \Theta_{m,j}|\{\sigma^j,\psi_m\}\rangle\langle\{\sigma^j,\psi_m\}|, \tag{73}$$

$$\Theta_{m,j} \equiv \frac{\Delta}{2}\sigma^j\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_0-z_{-1})L_+-(\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle + \frac{\bar{\Delta}}{2}\sigma^{-j}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_0-z_{-1})L_+-(\bar{z}_0-\bar{z}_{-1})L_-\}}|m\rangle\rangle. \tag{74}$$

Now for the latter use let us determine $\Theta_{m,j}$:

$$\Theta_{m,j} = \begin{cases}
(N) \quad \frac{1}{2}\left(\Delta\sigma^j + \bar{\Delta}\sigma^{-j}\right)\exp\left(-\frac{1}{2}|\kappa|^2\right)L_m\left(|\kappa|^2\right) \\
\qquad\qquad\qquad\qquad \text{where} \quad \kappa = C\sin(\pi/n)i, \\
(K) \quad \frac{1}{2}\left(\Delta\sigma^j + \bar{\Delta}\sigma^{-j}\right)\frac{m!}{(2K)_m}(1+|\kappa|^2)^{-K-m}F_m(|\kappa|^2 : 2K) \\
\qquad\qquad\qquad\qquad \text{where} \quad \kappa = \sinh\left(C\sin(\pi/n)\right)i, \\
(J) \quad \frac{1}{2}\left(\Delta\sigma^j + \bar{\Delta}\sigma^{-j}\right)\frac{m!}{{}_{2J}P_m}(1-|\kappa|^2)^{J-m}F_m(|\kappa|^2 : 2J) \\
\qquad\qquad\qquad\qquad \text{where} \quad \kappa = \sin\left(C\sin(\pi/n)\right)i,
\end{cases} \tag{75}$$

where C is a constant defined in (67).

Therefore let us solve (60)

$$i\frac{d}{dt}\Psi_0=\left(H_F+{H_F}^{\dagger}\right)\Psi_0=\left\{\left({H_F}^{'}+\tilde{H}_F^{'}\right)+\left({H_F}^{''}+\tilde{H}_F^{''}\right)\right\}\Psi_0 \tag{76}$$

with ${H_F}^{'}+\tilde{H}_F^{'}=\sum_m\sum_j\Theta_{m,j}|\{\sigma^j,\psi_m\}\rangle\langle\{\sigma^j,\psi_m\}|$. By making use of the method of constant variation again we can set $\Psi_0(t)$ as

$$\Psi_0(t)=\sum_{m=0}^{\infty}\sum_{j=0}^{n-1}\mathrm{e}^{-it\Theta_{m,j}}a_{m,j}(t)|\{\sigma^j,\psi_m\}\rangle, \tag{77}$$

then we have a set of extremely complicated equations with respect to $\{a_{m,j}\}$. It is almost impossible to solve them. Therefore we consider a simple case : for $m<r$

$$\Psi_0(t)=\sum_{j=0}^{n-1}\mathrm{e}^{-it\Theta_{m,j}}a_{m,j}(t)|\{\sigma^j,\psi_m\}\rangle+\sum_{j=0}^{n-1}\mathrm{e}^{-it\Theta_{r,j}}a_{r,j}(t)|\{\sigma^j,\psi_r\}\rangle. \tag{78}$$

That is, we adopt only two terms with respect to $\{m|m\geq 0\}$. Some careful algebras lead us to $(0\leq j\leq n-1)$

$$\begin{aligned}
&i\frac{d}{dt}a_{m,j}=\\
&\sum_{j'=0}^{n-1}\mathrm{e}^{it\Omega(m-r)}\mathrm{e}^{it(\Theta_{m,j}-\Theta_{r,j'})}\left[\frac{\Delta}{2}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\frac{\sigma^{kj-(k-1)j'}}{n}\right\}\right.\\
&\qquad\left.+\frac{\bar{\Delta}}{2}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\frac{\sigma^{(k-1)j-kj'}}{n}\right\}\right]a_{r,j'}\ ,
\end{aligned} \tag{79}$$

$$\begin{aligned}
&i\frac{d}{dt}a_{r,j}=\\
&\sum_{j'=0}^{n-1}\mathrm{e}^{it\Omega(r-m)}\mathrm{e}^{it(\Theta_{r,j}-\Theta_{m,j'})}\left[\frac{\Delta}{2}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\frac{\sigma^{kj-(k-1)j'}}{n}\right\}\right.\\
&\qquad\left.+\frac{\bar{\Delta}}{2}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\frac{\sigma^{(k-1)j-kj'}}{n}\right\}\right]a_{m,j'}\ .
\end{aligned}$$

However we cannot still solve the above equations exactly, so let us make the **rotating–wave approximation** from this stage. This means that the resonance condition is

$$\Omega(m-r)+(\Theta_{m,j}-\Theta_{r,j'})=0 \tag{80}$$

for some j and $j^{'}$, and we reject the remaining terms in (79).

A comment is in order. This resonance condition makes some constraint on the parameters $\{\omega, g, \Delta, \bar{\Delta}\}$ contained in the Hamiltonians (46), so ω, g and Δ, $\bar{\Delta}$ are already not free parameters.

As for a deep meaning of this approximation see [35], [36].

Then we obtain relatively simple equations :

$$\begin{aligned}
i\frac{d}{dt}a_{m,j} = &\left[\frac{\Delta}{2}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\frac{\sigma^{kj-(k-1)j^{'}}}{n}\right\}\right.\\
&\left.+\frac{\bar{\Delta}}{2}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\frac{\sigma^{(k-1)j-kj^{'}}}{n}\right\}\right]a_{r,j^{'}}\ ,\\
i\frac{d}{dt}a_{r,j^{'}} = &\left[\frac{\Delta}{2}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\frac{\sigma^{kj^{'}-(k-1)j}}{n}\right\}\right.\\
&\left.+\frac{\bar{\Delta}}{2}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\frac{\sigma^{(k-1)j^{'}-kj}}{n}\right\}\right]a_{m,j}\ .
\end{aligned}\tag{81}$$

Summing up the contents, we have

General Case ($j = j^{'}$ and $j \neq j^{'}$) $\qquad \Omega(m-r)+(\Theta_{m,j}-\Theta_{r,j^{'}})=0$

$$\begin{aligned}
i\frac{d}{dt}a_{m,j} = \frac{1}{2}&\left[\Delta\frac{\sigma^{j^{'}}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\sigma^{k(j-j^{'})}\right\}\right.\\
&\left.+\bar{\Delta}\frac{\sigma^{-j}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\sigma^{k(j-j^{'})}\right\}\right]a_{r,j^{'}}\ ,\\
i\frac{d}{dt}a_{r,j^{'}} = \frac{1}{2}&\left[\Delta\frac{\sigma^{j}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\sigma^{k(j^{'}-j)}\right\}\right.\\
&\left.+\bar{\Delta}\frac{\sigma^{-j^{'}}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\sigma^{k(j^{'}-j)}\right\}\right]a_{m,j}\ .
\end{aligned}\tag{82}$$

Here if we define the coefficient in the equations above as

$$\begin{aligned}
\mathcal{R}_{j^{'},j} =&\Delta\frac{\sigma^{j}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\sigma^{k(j^{'}-j)}\right\}\\
&+\bar{\Delta}\frac{\sigma^{-j^{'}}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle\sigma^{k(j^{'}-j)}\right\},
\end{aligned}\tag{83}$$

then

$$\begin{aligned}\bar{\mathcal{R}}_{j',j} =&\bar{\Delta}\frac{\sigma^{-j}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\sigma^{k(j-j^{'})}\right\} \\ &+\Delta\frac{\sigma^{j^{'}}}{n}\left\{\sum_{k=0}^{n-1}\langle\langle m|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|r\rangle\rangle\sigma^{k(j-j^{'})}\right\}.\end{aligned} \tag{84}$$

To determine $\mathcal{R}_{j',j}$ let us calculate a term $\langle\langle r|\mathrm{e}^{-\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle$:

$$\langle\langle r|\mathrm{e}^{\frac{1}{2}\{(z_k-z_{k-1})L_+-(\bar{z}_k-\bar{z}_{k-1})L_-\}}|m\rangle\rangle$$

$$= \begin{cases} (N) \quad \sqrt{\frac{m!}{r!}}\exp\left(-\frac{1}{2}|\kappa|^2\right)L_m{}^{(r-m)}\left(|\kappa|^2\right)\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sigma^{k(r-m)} \\ \qquad\qquad\qquad\qquad \text{where} \quad \kappa = C\mathrm{sin}(\pi/n)i, \\ (K) \quad \sqrt{\frac{r!m!}{(2K)_r(2K)_m}}(1+|\kappa|^2)^{-K-(r+m)/2}F_m{}^{(r-m)}(|\kappa|^2:2K)\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sigma^{k(r-m)} \\ \qquad\qquad\qquad\qquad \text{where} \quad \kappa = \sinh\left(C\mathrm{sin}(\pi/n)\right)i, \\ (J) \quad \sqrt{\frac{r!m!}{{}_{2J}P_r{}_{2J}P_m}}(1-|\kappa|^2)^{J-(r+m)/2}F_m{}^{(r-m)}(|\kappa|^2:2J)\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sigma^{k(r-m)} \\ \qquad\qquad\qquad\qquad \text{where} \quad \kappa = \sin\left(C\mathrm{sin}(\pi/n)\right)i, \end{cases} \tag{85}$$

where C is a constant defined in (67). 8

From this we obtain respectively

$$\begin{aligned}\mathcal{R}_{j',j} =&\left\{\Delta\frac{\sigma^{j}}{n}+\bar{\Delta}\frac{\sigma^{-j^{'}}}{n}(-1)^{r-m}\right\}\sqrt{\frac{m!}{r!}}\exp\left(-\frac{1}{2}|\kappa|^2\right)L_m{}^{(r-m)}\left(|\kappa|^2\right)\times \\ &\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sum_{k=0}^{n-1}\sigma^{k(r-m+j^{'}-j)},\end{aligned} \tag{86}$$

$$\begin{aligned}\mathcal{R}_{j',j} =&\left\{\Delta\frac{\sigma^{j}}{n}+\bar{\Delta}\frac{\sigma^{-j^{'}}}{n}(-1)^{r-m}\right\}\sqrt{\frac{r!m!}{(2K)_r(2K)_m}}(1+|\kappa|^2)^{-K-(r+m)/2}F_m{}^{(r-m)}(|\kappa|^2:2K)\times \\ &\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sum_{k=0}^{n-1}\sigma^{k(r-m+j^{'}-j)},\end{aligned} \tag{87}$$

$$\begin{aligned}\mathcal{R}_{j',j} =&\left\{\Delta\frac{\sigma^{j}}{n}+\bar{\Delta}\frac{\sigma^{-j^{'}}}{n}(-1)^{r-m}\right\}\sqrt{\frac{r!m!}{{}_{2J}P_r{}_{2J}P_m}}(1-|\kappa|^2)^{J-(r+m)/2}F_m{}^{(r-m)}(|\kappa|^2:2J)\times \\ &\kappa^{r-m}\sigma^{-(1/2)(r-m)}\sum_{k=0}^{n-1}\sigma^{k(r-m+j^{'}-j)}.\end{aligned} \tag{88}$$

Here it is easy to see

$$\sum_{k=0}^{n-1} \sigma^{k(r-m+j^{'}-j)} = \begin{cases} 0 & r-m+j^{'}-j \neq 0 \ (\mathrm{mod}\ n), \\ n & r-m+j^{'}-j = 0 \ (\mathrm{mod}\ n), \end{cases}$$

so we have only to consider the case $\quad r-m+j^{'}-j = 0 \ (\mathrm{mod}\ n) \quad$ in the following.

Now we are in a position to solve (82). For simplicity we set $\mathcal{R} = \mathcal{R}_{j^{'},j}$, then

$$i\frac{d}{dt}\begin{pmatrix} a_{m,j} \\ a_{r,j^{'}} \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{2}\bar{\mathcal{R}} \\ \frac{1}{2}\mathcal{R} & 0 \end{pmatrix}\begin{pmatrix} a_{m,j} \\ a_{r,j^{'}} \end{pmatrix}, \tag{89}$$

so their solutions are given by

$$\begin{pmatrix} a_{m,j}(t) \\ a_{r,j^{'}}(t) \end{pmatrix} = \begin{pmatrix} \cos(\frac{|\mathcal{R}|}{2}t) & -i\frac{\bar{\mathcal{R}}}{|\mathcal{R}|}\sin(\frac{|\mathcal{R}|}{2}t) \\ -i\frac{\mathcal{R}}{|\mathcal{R}|}\sin(\frac{|\mathcal{R}|}{2}t) & \cos(\frac{|\mathcal{R}|}{2}t) \end{pmatrix}\begin{pmatrix} a_{m,j}(0) \\ a_{n,j^{'}}(0) \end{pmatrix}. \tag{90}$$

That is, $\mathcal{R}$'s are just the (complex) Rabi frequencies. As shown in (86), (87), (88) they are very complicated.

A comment is in order. If we write $\mathcal{R} = z = \mathrm{e}^{i\theta}r$, then the above matrix becomes

$$\begin{pmatrix} \cos(\frac{r}{2}t) & -i\mathrm{e}^{-i\theta}\sin(\frac{r}{2}t) \\ -i\mathrm{e}^{i\theta}\sin(\frac{r}{2}t) & \cos(\frac{r}{2}t) \end{pmatrix}. \tag{91}$$

This is a form required in Quantum Computation. However in the 2–level case $\mathcal{R}$ is real, so the matrix becomes

$$\begin{pmatrix} \cos(\frac{r}{2}t) & -i\sin(\frac{r}{2}t) \\ -i\sin(\frac{r}{2}t) & \cos(\frac{r}{2}t) \end{pmatrix},$$

see [17]. This is too special to perform Quantum Computation. Our result is a bonus due to the generalization from 2–level system to multi level one.

We have verified the Rabi oscillations in our extended models under the rotating–wave approximation, so there are many things to be performed from Quantum Optics or Mathematical Physics. In the forthcoming papers we will report them.

Here let us present an (good ?) exercise to the readers :

Exercise For $m < r < s$ we set

$$
\begin{aligned}
&\Psi_0(t) \\
=&\sum_{j=0}^{n-1} \mathrm{e}^{-it\Theta_{m,j}} a_{m,j}(t)|\{\sigma^j, \psi_m\}\rangle + \sum_{j=0}^{n-1} \mathrm{e}^{-it\Theta_{r,j}} a_{r,j}(t)|\{\sigma^j, \psi_r\}\rangle + \sum_{j=0}^{n-1} \mathrm{e}^{-it\Theta_{s,j}} a_{s,j}(t)|\{\sigma^j, \psi_s\}\rangle.
\end{aligned}
$$

Solve the equation (76) under this ansatz. Then you must pay attention to the consistency of some resonance conditions like (80).

5 Unitary Operations in Quantum Computation

In this section we make a brief comment on application of the results in the preceding section to quantum computation based on qudits space. Remind once more that the following arguments are based on the **rotating wave approximation**.

First we must solve the equation which j and $j^{'}$ satisfy

$$
j^{'} - j + r - m = 0 \ (\mathrm{mod}\ n) \quad \text{for} \quad m < r. \tag{92}
$$

The solutions are easily given as follows.

$$
\begin{aligned}
(j^{'},\ j) = {} & (n - r + m,\ 0),\ (n - r + m + 1,\ 1),\ \cdots,\ (n - 1,\ r - m - 1), \\
& (n - r + m - 1,\ n - 1),\ (n - r + m - 2,\ n - 2),\ \cdots,\ (0,\ r - m).
\end{aligned} \tag{93}
$$

Here let us rewrite the equation (89) in a full form

$$
i\frac{d}{dt}
\begin{pmatrix}
a_{m,0} \\ \cdot \\ a_{m,j} \\ \cdot \\ a_{m,n-1} \\ a_{r,0} \\ \cdot \\ a_{r,j'} \\ \cdot \\ a_{r,n-1}
\end{pmatrix}
=
\begin{pmatrix}
0 & & & & & & & & & \\
 & \cdot & & & & & & & & \\
 & & 0 & & & & & \frac{1}{2}\bar{\mathcal{R}} & & \\
 & & & \cdot & & & & & & \\
 & & & & \cdot & & & & & \\
 & & & & & \cdot & & & & \\
 & & & & & & \cdot & & & \\
 & & \frac{1}{2}\mathcal{R} & & & & & 0 & & \\
 & & & & & & & & \cdot & \\
 & & & & & & & & & 0
\end{pmatrix}
\begin{pmatrix}
a_{m,0} \\ \cdot \\ a_{m,j} \\ \cdot \\ a_{m,n-1} \\ a_{r,0} \\ \cdot \\ a_{r,j'} \\ \cdot \\ a_{r,n-1}
\end{pmatrix}
\tag{94}
$$

where $\mathcal{R} = \mathcal{R}_{j',j}$. Then the solution (corresponding to (90)) is explicitly

$$
\begin{pmatrix}
a_{m,0}(t) \\ \cdot \\ a_{m,j}(t) \\ \cdot \\ a_{m,n-1}(t) \\ a_{r,0}(t) \\ \cdot \\ a_{r,j'}(t) \\ \cdot \\ a_{r,n-1}(t)
\end{pmatrix}
=
\begin{pmatrix}
1 & & & & & & & & & \\
& \cdot & & & & & & & & \\
& & \cos(\frac{|\mathcal{R}|}{2}t) & & & & & -i\frac{\bar{\mathcal{R}}}{|\mathcal{R}|}\sin(\frac{|\mathcal{R}|}{2}t) & & \\
& & & \cdot & & & & & & \\
& & & & \cdot & & & & & \\
& & & & & \cdot & & & & \\
& & & & & & \cdot & & & \\
& & -i\frac{\mathcal{R}}{|\mathcal{R}|}\sin(\frac{|\mathcal{R}|}{2}t) & & & & & \cos(\frac{|\mathcal{R}|}{2}t) & & \\
& & & & & & & & \cdot & \\
& & & & & & & & & 1
\end{pmatrix}
\begin{pmatrix}
a_{m,0}(0) \\ \cdot \\ a_{m,j}(0) \\ \cdot \\ a_{m,n-1}(0) \\ a_{r,0}(0) \\ \cdot \\ a_{r,j'}(0) \\ \cdot \\ a_{r,n-1}(0)
\end{pmatrix}. \tag{95}
$$

That is, we have a unitary operation which will be written as $U(j, j^{'}; t)$. This operation is crucial in Quantum Computation.

As a result we obtained n elementary unitary operations

$$
\begin{aligned}
&U(0, n-r+m; t),\ U(1, n-r+m+1; t),\ \cdots,\ U(r-m-1, n-1; t), \\
&U(r-m, 0; t),\ U(r-m+1, 1; t),\ \cdots,\ U(n-1, n-r+m-1; t).
\end{aligned} \tag{96}
$$

One can combine these operations suitably to obtain the (complex) operation required in

Quantum Computation.

We would like to apply these operations to Quantum Computation on the space of 2–qudits. For that we need some modification in the preceeding section. Namely, we must consider a set of n states : for $m_0 < m_1 < \cdots < m_{n-1}$

$$\Psi_0(t) = \sum_{k=0}^{n-1} \sum_{j=0}^{n-1} \mathrm{e}^{-it\Theta_{m(k),j}} a_{m(k),j}(t) |\{\sigma^j, \psi_{m(k)}\}\rangle \tag{97}$$

instead of (78). Of course for any pair $m_k < m_l$ we have the same result. We cannot solve the equation at a stroke, so we reduce the calculation on n–system to that of the pairs. Remind once more that in this paper we are interested in Rabi oscillations between two states. By combining these Rabi oscillations one after another we obtain unitary operations in our full system.

For (97) we have the huge (block-) equation

$$\begin{pmatrix} \mathbf{a_{m(0),j}}(t) \\ \mathbf{a_{m(1),j}}(t) \\ \cdot \\ \cdot \\ \cdot \\ \mathbf{a_{m(n-1),j}}(t) \end{pmatrix} = \begin{pmatrix} A_{00} & A_{01} & \cdot\ \cdot\ \cdot & A_{0,n-1} \\ A_{10} & A_{11} & \cdot\ \cdot\ \cdot & A_{1,n-1} \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ A_{n-1,0} & A_{n-1,1} & \cdot\ \cdot\ \cdot & A_{n-1,n-1} \end{pmatrix} \begin{pmatrix} \mathbf{a_{m(0),j}}(0) \\ \mathbf{a_{m(1),j}}(0) \\ \cdot \\ \cdot \\ \cdot \\ \mathbf{a_{m(n-1),j}}(0) \end{pmatrix} \tag{98}$$

where each $\mathbf{a_{m(k),j}}$ is a n vector and A_{ij} a $n \times\ n$ matrix.

For example, if we pay attention to the pair $m_k < m_l$ then the block (in the above huge matrix)

$$\begin{pmatrix} A_{m(k)m(k)} & A_{m(k)m(l)} \\ A_{m(l)m(k)} & A_{m(l)m(l)} \end{pmatrix} \tag{99}$$

is just one of above unitary operations in (96) $(m = m_k,\ r = m_l)$ and the remainders are $A_{pp} = \mathbf{1}$, $A_{pq} = \mathbf{0}$ $(p \neq q)$. As a total we have the elementary unitary transformations in $U(n^2)$ with the following number

$$n\,{}_nC_2 = \frac{n^2(n-1)}{2}.$$

Here if we identify (97) with an element on the space of 2–qudits like

$$\sum_{k=0}^{n-1}\sum_{j=0}^{n-1} a_{k,j}(t)|k\rangle \otimes |j\rangle \ \in\ \mathbf{C}^n \otimes \mathbf{C}^n \tag{100}$$

where $\mathbf{C}^n = \mathrm{Vect}_{\mathbf{C}}\{|0\rangle, |1\rangle, \cdots, |n-1\rangle\}$, then we have plenty of unitary transformations in $U(n^2)$. We want to construct or perform some logic gates on the space of 2–qudits, see [24], [25] for details.

For example, we would like to construct the (reverse) controlled–shift gate ((reverse) controlled–unitary ones more generally) by combining elementary unitary operations obtained in this paper :



where Σ in the above figure is just Σ_1 in (39), and a and b run from 0 to $n-1$.

We would like to conclude this section by stating that a possible realization of our model could be found in Josephson junctions [11] and ion cavities [37], [38], [39].

6 Discussion

In this paper we extended the works in [12], [14] and [17] based on 2–level system to n–level system and obtained the general theory of Rabi frequencies in the strong coupling regime under the rotating–wave approximation. The essence of our work is that Rabi frequencies are in this case deeply connected to the matrix elements of coherent and/or generalized coherent operators based on $su(2)$ and $su(1,1)$, [16], [8].

One of motivations of this study is to apply our results to Holonomic Quantum Computation developed by Italian group (Pachos, Rasetti and Zanardi) and the author, see [26], [27], [28], [29] and [30]—[34].

Holonomic Quantum Computation is based on 2–level system at this stage, but it is very natural to extend it from 2–level system to n–level system (let us call it Generalized Holonomic Quantum Computation (GHQC)). In fact the author has attempted its extension in [30], but it was not complete due to some lack of new method.

As shown in this paper we have obtained some new method or technique, so we would like to challenge once more.

Acknowledgment.
The author wishes to thank Marco Frasca and Zakaria Giunashvili for their helpful comments and suggestions.

References

[1] J. R. Klauder and Bo-S. Skagerstam (Eds) : Coherent States, World Scientific, Singapore, 1985.

[2] L. Mandel and E. Wolf : Optical Coherence and Quantum Optics, Cambridge University Press, 1995.

[3] A. Perelomov : Generalized Coherent States and Their Applications, Springer–Verlag, 1986.

[4] K. Funahashi, T. Kashiwa, S. Sakoda and K. Fujii : Coherent states, path integral, and semiclassical approximation, J. Math. Phys., 36(1995), 3232.

[5] K. Funahashi, T. Kashiwa, S. Sakoda and K. Fujii : Exactness in the Wentzel-Kramers-Brillouin approximation for some homogeneous spaces, J. Math. Phys., 36(1995), 4590.

[6] K. Fujii, T. Kashiwa, S. Sakoda :Coherent states over Grassmann manifolds and the WKB exactness in path integral, J. Math. Phys., 37(1996), 567.

[7] K. Fujii : Basic Properties of Coherent and Generalized Coherent Operators Revisited, Mod. Phys. Lett. A 16(2001), 1277, quant-ph/0009012.

[8] K. Fujii : Note on Extended Coherent Operators and Some Basic Properties, submitted, quant-ph/0009116.

[9] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, to appear in J. Applied Math, quant-ph/0103011.

[10] P. Meystre and M. Sargent III : Elements of Quantum Optics, Springer–Verlag, 1990–1991.

[11] Y. Nakamura, Yu. A. Pashkin and J. S. Tsai : Rabi Oscillations in a Josephson–Junction Charge Two–Level System, Phys. Rev. Lett., 87(2001), 246601.

[12] M. Frasca : Rabi oscillations and macroscopic quantum superposition states, Phys. Rev. A 66(2002), 023810, quant-ph/0111134.

[13] M. Frasca : Dephasing by two–level systems at zero temperature by unitary evolution, to appear in Physica E, cond-mat/0112253.

[14] M. Frasca : Classical states and decoherence produced by unitary evolution in the thermodynamic limit, J. Opt. B : Quantum Semiclass. Opt. 4(2002), S443, quant-ph/0201040.

[15] M. Frasca : A modern review of the two–level approximationn, quant-ph/0209056.

[16] K. Fujii : Matrix Elements of Generalized Coherent Operators, submitted, quant-ph/0202081.

[17] K. Fujii : Mathematical Structure of Rabi Oscillations in the Strong Coupling Regime, to appear in J. Phys. A, quant-ph/0203135.

[18] E. Tur : Jaynes-Cummings model without rotating wave approximation. asymptotics of eigenvalues, math-ph/0211055.

[19] W. P. Schleich : Quantum Optics in Phase Space, Wiley–VCH, 2001.

[20] A. Muthukrishnan and C. R. Stroud, Jr. : Multivalued logic gates for quantum computation, Phys. Rev. A 62(2000), 052309.

[21] N. J. Cerf, M. Bourennane, A. Karlsson and N. Gisin : Security of quantum key distribution using d–level systems, quant-ph/0107130.

[22] V. Karimipour, S. Bagherinezhad and A. Bahraminasab : Quantum Key Distribution for d–level systems with Generalized Bell States, quant-ph/0111091.

[23] M. A. Nielsen, M. J. Bremner and J. L. Dodd : Universal simulation of Hamiltonian dynamics for equation systems with finite–dimensional state spaces, quant-ph/0109064.

[24] K. Fujii : Exchange Gate on the Qudit Space and Fock Space, submitted, quant-ph/0207002.

[25] K. Funahashi : The Controlled–U and unitary transformations on two qudits, in preparation.

[26] P. Zanardi and M. Rasetti : Holonomic Quantum Computation, Phys. Lett. A 264(1999), 94, quant-ph/9904011.

[27] J. Pachos, P. Zanardi and M. Rasetti : Non-Abelian Berry connections for quantum computation, Phys. Rev. A 61(2000), 010305(R), quant-ph/9907103.

[28] J. Pachos and S. Chountasis : Optical Holonomic Quantum Computer, Phys. Rev. A 62(2000), 052318, quant-ph/9912093.

[29] J. Pachos and P. Zanardi : Quantum Holonomies for Quantum Computing, Int. J. Mod. Phys. B 15(2001), 1257, quant-ph/0007110.

[30] K. Fujii : Note on Coherent States and Adiabatic Connections, Curvatures, J. Math. Phys. 41(2000), 4406, quant-ph/9910069.

[31] K. Fujii : Mathematical Foundations of Holonomic Quantum Computer, Rept. Math. Phys. 48(2001), 75, quant-ph/0004102.

[32] K. Fujii : More on Optical Holonomic Quantum Computer, quant-ph/0005129.

[33] K. Fujii : Mathematical Foundations of Holonomic Quantum Computer II, quant-ph/00101102.

[34] K. Fujii : From Geometry to Quantum Computation, to appear in Proceedings of the 2nd International Symposium " Quantum Theory and Symmetries", World Scientific, quant-ph/0107128.

[35] M. Frasca : Theory of quantum resonance : A renormalization–group approach, Phys. Rev. A 58(1998), 771.

[36] S. Ei, K. Fujii and T. Kunihiro : Renormalization–Group Method for Reduction of Evolution Equations ; Invariant Manifolds and Envelopes, Ann. of Physics, 280(2000), 236, hep–th/9905088.

[37] C. Montoe, D. Leibfried, B. E. King, D. M. Meekhof, W. M. Itano and D. J. Wineland : Simplified Quantum Logic with Trapped Ions, quant-ph/9612042.

[38] W. M. Itano, C. Montoe, D. M. Meekhof, D. Leibfried, B. E. King and D. J. Wineland : Quantum harmonic oscillator state synthesis and analysis, quant-ph/9702038.

[39] H. Moya-Cessa, A. Vidiella, J. A. Roversi, D. S. Freitas and S. M. Dutra : Long–time–scale revivals in ion traps, Phys. Rev. A 59(1999), 2518.